Interaction-induced Bose Metal in 2D

Denis Dalidovich and Philip Phillips
Loomis Laboratory of Physics
University of Illinois at Urbana-Champaign
1100 W.Green St., Urbana, IL, 61801-3080

We show here that the regularization of the conductivity resulting from the bosonic interactions on the 'insulating' (quantum disordered) side of an insulator-superconductor transition in 2D gives rise to a metal with a finite conductivity, $\sigma = (2/\pi)4e^2/h$, as temperature tends to zero. The Bose metal is stable to weak disorder and hence represents a concrete example of an interaction-induced metallic phase. Phenomenological inclusion of dissipation reinstates the anticipated insulating behaviour in the quantum-disordered regime. Hence, we conclude that the traditionally-studied insulator-superconductor transition, which is driven solely by quantum fluctuations, corresponds to a superconductor-metal transition. The possible relationship to experiments on superconducting thin films in which a low-temperature metallic phase has been observed is discussed.

arXiv:cond-mat/0005119v3 29 May 2001

Two ground states are thought to exist for bosons at $T = 0$: a superconductor with long-range phase coherence and an insulator in which the quantum mechanical phase is disordered. In this paper, we prove that the phase-disordered regime is actually not an insulator but rather a metal with a universal conductivity given by $(2/\pi)4e^2/h$. This result arises from a conspiracy: In the quantum-disordered regime, the population of bosons is exponentially suppressed; however, so is the scattering rate between bosons. But because the conductivity is a product of the density and the scattering time, the exponentials cancel, giving rise to a finite conductivity at $T = 0$. Any finite amount of dissipation, however, reinstates the insulator. Consequently, the traditional arguments for the insulating phase of bosons based on quantum fluctuations of the phase must be reconsidered. Quantum fluctuations alone are insufficient to yield an insulating phase as illustrated clearly in Fig. (1). Generically, quantum fluctuations lead to a Bose metal phase in 2D.

To establish our result, we rely on the Landau-Ginzburg formalism. While this approach has had much success [1,2,4–12] in elucidating the critical properties of thin films and Josephson junction arrays at the point of transition [4,13], the transport properties have proven to be more elusive. For example, simple physical considerations [2,3] place the conductivity at the point of transition at a universal value of $\sigma_Q = 4e^2/h$, whereas in experiments, σ_Q ranges anywhere from $2\sigma_Q$ to $\sigma_Q/3$ [14–16]. Likewise, theoretical calculations based on the Landau-Ginsburg approach have yielded values ranging from $\pi\sigma_Q/8$ [11,12] to $1.037\sigma_Q$ [17]. In addition, Damle and Sachdev [17] have shown that the frequency and temperature tending to zero limits do not commute. The lack of commutativity of these two limits arises from the fundamental fact that close to the transition point, the conductivity is a universal function of $\hbar\omega/T$ [17]. Consequently, $\sigma(\omega \to 0, T = 0) \neq \sigma(\omega = 0, T \to 0)$. The original theoretical work [2,12] was all based on the former limit which physically represents the coherent regime.



FIG. 1. Phase diagram for the superconductor-Bose metal transition as a function of temperature and the quantum fluctuations, g. For Josephson-junction arrays, g is given by the ratio of the charging energy, E_c, to the Josephson coupling, J. $T_{\rm KT}$ is the Kosterlitz-Thouless temperature at which phase coherence obtains. g_c defines the critical value of the phase disorder to destroy the superconducting phase. QC refers to the quantum critical regime in which the inverse correlation length is linear in the temperature.

In such calculations, the dc-conductivity acquires a singular Drude part on the insulating side which can be regularized by the phenomenological inclusion of dissipation [11,18]. Experimentally, however, it is $\sigma(\omega = 0, T \to 0)$, or equivalently the hydrodynamic regime, that is relevant. In this regime, collisions between quasiparticles dominates the conductivity which can be obtained from the quantum kinetic equation. While such an approach has been used in the quantum critical

regime, no corresponding study has been made in the quantum-disordered regime. It is precisely this limit that we study here using the quantum kinetic approach. We show explicitly that the quartic interaction between the bosons, whose dispersion relation is gapped in the phase-disordered regime, ultimately gives rise to a metallic rather than the anticipated insulating phase in the quantum-disordered regime. The Bose metal that we find is distinct from the Bose metal of Das and Doniach [19] which is purported to obtain once the size of the Josephson superconducting grains exceeds a critical value. Further, our result establishes that the Bose metal is the generic ground state of bosons lacking phase coherence in the absence of dissipation.

Although our microscopic system is an array of Josephson junctions, we coarse-grain over the phase associated with each junction and hence use as our starting point the Landau-Ginzburg action,

$$F[\psi] = \sum_{\mathbf{k},\omega_n} (k^2 + \omega_n^2 + m^2)|\psi(\mathbf{k},\omega_n)|^2 + \frac{U}{2N\beta} \sum_{\omega_1,\ldots,\omega_4;\mathbf{k}_1,\ldots,\mathbf{k}_4} \delta_{\omega_1+\cdots\omega_4,0}\delta_{\mathbf{k}_1+\cdots\mathbf{k}_4,0} \psi_\nu(\omega_1,\mathbf{k}_1)\psi_\nu(\omega_2,\mathbf{k}_2)\psi_\mu(\omega_3,\mathbf{k}_3)\psi_\mu(\omega_4,\mathbf{k}_4) \quad (1)$$

where $\psi(\mathbf{r},\tau)$ is the complex Bose order parameter whose expectation value is proportional to $\langle \exp(i\phi)\rangle$, where ϕ is the phase of a particular junction. The summation in the action over discrete Matsubara frequencies, $\omega_i = 2\pi n_i T$, and integration over continuous wavevectors, $\mathbf{k}$, is assumed. The parameter m^2 is the inverse square of the correlation length. In writing the action in this fashion, we have already included the one-loop renormalization arising from the quartic term. In the quantum-disordered regime, $m \gg T$ and hence it is the quantum fluctuations that dominate the divergence of the correlation length.

Our goal is to calculate the conductivity in the quantum disordered regime in the collision-dominated limit. Following the quantum kinetic approach of Damle and Sachdev [17], we introduce the distribution function, $f(\mathbf{k},t)$ for quasiparticles. We have suppressed the distinction between particles and holes as they have identical distribution functions. The quantum kinetic equation for the quasiparticle distribution function $f(\mathbf{k},t)$ takes the following form

$$\frac{\partial}{\partial t}\delta f(\mathbf{k},t) + e^*\mathbf{E}(t)\frac{\partial}{\partial \mathbf{k}} n(\epsilon_{\mathbf{k}}) = I\{\delta f\} \quad (2)$$

when it is linearised in the correction $\delta f(\mathbf{k},t)$ to the equilibrium Bose distribution $n(\epsilon_k) = (e^{\epsilon_k/T}-1)^{-1}$ that is induced by the electric field $\mathbf{E}(t)$. The gapped dispersion relation for the bosons is $\epsilon_{\mathbf{k}} = \sqrt{k^2+m^2}$. The resultant linearised collision integral [20]

$$I\{\delta f\} = -\frac{1}{N(2\pi)^2}\left[\int_{\epsilon_{\mathbf{q}+\mathbf{k}}>\epsilon_{\mathbf{k}}} d^2q \frac{1}{\epsilon_{\mathbf{q}+\mathbf{k}}\epsilon_{\mathbf{k}}}\mathrm{Im}\frac{1}{\Pi(\mathbf{q},\epsilon_{\mathbf{q}+\mathbf{k}}-\epsilon_{\mathbf{k}})} \right.$$
$$[\delta f(\mathbf{k})(n(\epsilon_{\mathbf{q}+\mathbf{k}}-\epsilon_{\mathbf{k}}) - n(\epsilon_{\mathbf{q}})) - \delta f(\mathbf{q}+\mathbf{k})(1+n(\epsilon_{\mathbf{k}})+n(\epsilon_{\mathbf{q}+\mathbf{k}}-\epsilon_{\mathbf{k}}))]$$
$$+\int_{\epsilon_{\mathbf{q}+\mathbf{k}}<\epsilon_{\mathbf{k}}} d^2q \frac{1}{\epsilon_{\mathbf{q}+\mathbf{k}}\epsilon_{\mathbf{k}}}\mathrm{Im}\frac{1}{\Pi(\mathbf{q},\epsilon_{\mathbf{k}}-\epsilon_{\mathbf{q}+\mathbf{k}})} \times [\delta f(\mathbf{k})(1+n(\epsilon_{\mathbf{q}})+n(\epsilon_{\mathbf{k}}-\epsilon_{\mathbf{q}+\mathbf{k}})) + \delta f(\mathbf{q}+\mathbf{k})(n(\epsilon_{\mathbf{k}})-n(\epsilon_{\mathbf{k}}-\epsilon_{\mathbf{q}+\mathbf{k}}))]$$
$$+\int_q d^2q \frac{1}{\epsilon_{\mathbf{q}}\epsilon_{\mathbf{k}}}\mathrm{Im}\frac{1}{\Pi(\mathbf{q}+\mathbf{k},\epsilon_{\mathbf{q}}+\epsilon_{\mathbf{k}})} [\delta f(\mathbf{k})(n(\epsilon_{\mathbf{q}})-n(\epsilon_{\mathbf{k}}+\epsilon_{\mathbf{q}})) \left. + \delta f(\mathbf{q})(n(\epsilon_{\mathbf{k}})-n(\epsilon_{\mathbf{k}}+\epsilon_{\mathbf{q}}))]\right] \quad (3)$$

is a sum of all incoming and outgoing quasiparticle scattering processes. A central quantity appearing in the collision integral is the polarization function

$$\Pi(\mathbf{q},i\Omega_m) = T\sum_n \int \frac{d^2p}{(2\pi)^2} G_0(\mathbf{p}+\mathbf{q},\omega_n+\Omega_m)G_0(\mathbf{p},\omega_n) \quad (4)$$

where the bare field propagator $G_0(\mathbf{p},\omega_n) = (p^2+\omega_n^2+m^2)^{-1}$. As it is the imaginary part of Π that is required in the collision integral, we must perform an analytical continuation $\Omega_n \to -i\Omega_n - \delta$ with δ a positive infinitesimal. Unlike Damle and Sachdev [17], we have found it convenient to work directly in d=2 within a large N expansion. Eq. (2) represents the linear integral equation that must be solved in the generic case to determine the role of the quartic term on the collision-induced conductivity.

To compute the conductivity, we work in the relaxation-time approximation in which $I\{\delta f\} \approx -\delta f(\mathbf{k})/\tau_{\mathbf{k}}$. In this approximation, the conductivity emerges as a momentum integration of the form,

$$\sigma = 2\frac{(e^*)^2}{\hbar}\int \frac{d^2k}{(2\pi)^2}\frac{k_x^2}{\epsilon_{\mathbf{k}}^2}\tau_{\mathbf{k}}\left(-\frac{\partial n(\epsilon_{\mathbf{k}})}{\partial \epsilon_{\mathbf{k}}}\right). \quad (5)$$

We must extract then the relaxation time, $\tau_{\mathbf{k}}$ from the collision integral. In the quantum-disordered regime, the statistics of the quasiparticles becomes Boltzmannian because $m \gg T$. This property results in a suppression of each subsequent $1/N$ correction by a factor $e^{-m/T}$ [21]. Having set $N=2$, we obtain then from Eq. (3) that

$$\frac{1}{\tau_{\mathbf{k}}} = \frac{1}{2(2\pi)^2}\left[\int \frac{d^2q}{\epsilon_{\mathbf{q}+\mathbf{k}}\epsilon_{\mathbf{k}}}\left(\mathrm{Im}\frac{1}{\Pi(\mathbf{q},\epsilon_{\mathbf{q}+\mathbf{k}}-\epsilon_{\mathbf{k}})}\right) n(\epsilon_{\mathbf{q}+\mathbf{k}}-\epsilon_{\mathbf{k}}) + \int \frac{d^2q}{\epsilon_{\mathbf{q}}\epsilon_{\mathbf{k}}}\left(\mathrm{Im}\frac{1}{\Pi(\mathbf{q}+\mathbf{k},\epsilon_{\mathbf{q}}+\epsilon_{\mathbf{k}})}\right) n(\epsilon_{\mathbf{q}})\right] \quad (6)$$

to leading accuracy. Here we took into account the fact that from Eq. (4), $\mathrm{Im}\Pi(\mathbf{q},\Omega)$ is an odd function of Ω and used the identity $1+n(\epsilon_{\mathbf{k}}-\epsilon_{\mathbf{q}+\mathbf{k}}) = -n(\epsilon_{\mathbf{q}+\mathbf{k}}-\epsilon_{\mathbf{k}})$.

The essence of our central result is that to leading order in T/m, the inverse relaxation time $1/\tau_{\mathbf{k}}$ is momentum-independent and given by

$$\frac{1}{\tau} = \pi T e^{-m/T}. \tag{7}$$

Substitution of this expression into Eq. (5) leads to the mutual cancellation of the exponential factors yielding the remarkable result

$$\sigma(T \to 0) = \frac{2}{\pi}\frac{4e^2}{h}. \tag{8}$$

It is curious to note [22] that a similar cancellation of exponential factors (from the mean free path and the density of states) arises in the context of the quasiparticle conductivity in a dirty d-wave superconductivity yielding the identical numerical prefactor $2/\pi$.

To establish this result, we obtain a workable expression for the inverse polarization function. In the leading approximation in T/m, the real part of the polarization function can be found from its value at $T = 0$ [21]:

$$\Pi(\mathbf{q}, \Omega) = \frac{1}{4\pi\sqrt{q^2 - \Omega^2}} \arctan \frac{\sqrt{q^2 - \Omega^2}}{2m} \tag{9}$$

It is easy to show from the above expression that when $\sqrt{\Omega^2 - q^2} > 2m$, the leading term of the imaginary part is temperature independent and equal to

$$\mathrm{Im}\Pi(\mathbf{q}, \Omega) = -\frac{1}{8\sqrt{\Omega^2 - q^2}}, \tag{10}$$

while in all other cases, such a contribution is absent and the leading temperature dependence comes from the first term in

$$\begin{aligned}\mathrm{Im}\Pi(\mathbf{q}, \Omega) = -\frac{1}{16\pi}\int \frac{d^2p}{\epsilon_{\mathbf{p}}\epsilon_{\mathbf{p+q}}}\{&|n(\epsilon_{\mathbf{p+q}}) - n(\epsilon_{\mathbf{p}})| \\ &\delta(|\epsilon_{\mathbf{p+q}} - \epsilon_{\mathbf{p}}| - \Omega) + (n(\epsilon_{\mathbf{p+q}}) + n(\epsilon_{\mathbf{p}})) \\ &\delta(\epsilon_{\mathbf{p+q}} + \epsilon_{\mathbf{p}} - \Omega)\},\end{aligned} \tag{11}$$

which is obtained by applying the Poisson summation formula to Eq. (4) [21]. The $\delta-$ functions are eliminated upon an angular integration. As we are concerned with the quantum-disordered regime, we focus on the limits $q, k \leq \sqrt{mT} \ll m$. Because in the first term in Eq. (6), $q > |\epsilon_{\mathbf{q+k}} - \epsilon_{\mathbf{k}}|$, we must use Eq. (11) to determine the imaginary part of the inverse polarization function. Using the fact that in the region of momentum integration, $|\mathrm{Re}\Pi(\mathbf{q}, \Omega)| \gg |\mathrm{Im}\Pi(\mathbf{q}, \Omega)|$, we obtain that

$$\begin{aligned}\mathrm{Im}\frac{1}{\Pi(\mathbf{q}, \epsilon_{\mathbf{q+k}} - \epsilon_{\mathbf{k}})} \approx \frac{8\pi m^2}{q}\sqrt{\frac{\pi T}{2m}} e^{-m/T} \cdot e^{-(q+(\mathbf{qk})/q)^2/2mT} \\ \left(e^{(\epsilon_{\mathbf{q+k}} - \epsilon_{\mathbf{k}})/T} - 1\right).\end{aligned} \tag{12}$$

In the second term of the Eq. (6) we have that $[(\epsilon_{\mathbf{q}} + \epsilon_{\mathbf{k}})^2 - (\mathbf{q} + \mathbf{k})^2] > 2m$. So, it would be sufficient to use Eq. (10) for the imaginary part, and the real part can be obtained from Eq. (9) yielding

$$\mathrm{Im}\frac{1}{\Pi(\mathbf{q} + \mathbf{k}, \epsilon_{\mathbf{q}} + \epsilon_{\mathbf{k}})} \approx \frac{16m}{\frac{4}{\pi^2}\ln^2\frac{4m}{|\mathbf{q}-\mathbf{k}|} + 1}. \tag{13}$$

This suggests that in the limit $m/T \to \infty$, the contribution to the relaxation time arising from the second term is logarithmically suppressed compared to the contribution from the first term. Substitution of Eq. (12) into Eq. (6) and performing the $q-$integration we arrive at the advertised result, Eq. (7), to leading accuracy in T/m.

This result is truly remarkable, because, as one can see, the conductivity in the quantum disordered regime (in the leading approximation) depends neither on temperature nor on the distance from the transition point. This means that the dc conductivity in this model, $\sigma = \sigma_Q g(m/T)$, where g is a universal function close to the transition point, displays a crossover upon lowering T from the universal value in the quantum critical regime [17], σ_Q, to the smaller value, Eq. (8) in the quantum-disordered regime. Though we focused on the relaxation time approximation, it is possible to verify *a posteriori* that the incoming term in the collision integral provides a contribution subdominant in T/m. Indeed, seeking the solution of the kinetic equation by means of consecutive approximations $\delta f(\mathbf{k}) = \delta f^{(0)}(\mathbf{k}) + \delta f^{(1)}(\mathbf{k}) + \cdots$, where

$$\delta f^{(0)}(\mathbf{k}) = -e^* \tau_{\mathbf{k}} \frac{(\mathbf{Ek})}{\epsilon_{\mathbf{k}}}\left(\frac{\partial n}{\partial \epsilon_{\mathbf{k}}}\right), \tag{14}$$

we obtain after substitution into the kinetic equation, that $\delta f^{(1)}(\mathbf{k})$ is proportional to higher powers of T/m than are the outgoing terms. Consequently, this subdominance justifies the relaxation time approximation in which only outgoing terms are retained.

The universal value for the conductivity, Eq. (8), was obtained for the region near the zero- temperature IST point, where $m \ll 1$. However, it is not difficult to see that the above results can be generalized for the case $m \approx O(1)$, keeping $T \ll 1$. One needs only to make the substitution in the collision integral

$$\frac{1}{N}\mathrm{Im}\frac{1}{\Pi(\mathbf{q}, \Omega)} \to \mathrm{Im}\frac{U}{1 + UN\Pi(\mathbf{q}, \Omega)}. \tag{15}$$

The subsequent steps are identical to those described above and yield the value of the conductivity

$$\sigma(T \to 0) = \frac{2}{\pi}\frac{U^2}{(4\pi m + U)^2}\frac{4e^2}{h}. \tag{16}$$

As expected, this value is not universal, and both m and U are the functions of parameters of the initial microscopic Hamiltonian. So, the leveled resistivity, generally speaking, depends on the distance from the transition point. In the limit $m \ll 1$, Eq. (16) reduces to the universal value (8). Note this result is specific to 2D. Because the conductivity is not [27] universal for the 3D system, the 3D case deserves special attention.

In the approximation that the inverse quasiparticle scattering rate is small relative to its energy, the general form for the conductivity, Eq. (5), can be obtained from the standard Kubo formula [11,18]. With this realization, the central result, Eq. (8), can be obtained from Eq. (8) of Ref. (18) by making the substitution $\eta \to 1/\tau$ ($\kappa = 1$), with $1/\tau$ given by Eq. (7).

It is worth exploring how robust our bose metal is as it has not yet been observed generically in experiments [24–26]. For s-wave pairing, the pair amplitude survives in the presence of weak disorder [28]. Consequently, the Bose metal remains intact in this limit. As a result, we have constructed a concrete example of an interaction-induced metal that persists in the limit of weak disorder. This is important as metallic phases have been observed in thin films [16,25,31] which should nominally exhibit only insulating or superconducting phases. For example, Jaeger, et. al. [31] have observed a downturn followed by a leveling of the resistance in Ga and In thin films with a saturation value ranging between $0.5k\Omega - 50k\Omega$. Our value of $(\pi/2)h/4e^2 \approx 10k\Omega$ is certainly within the experimental range over which the Bose metal phase has been observed. Hence the Bose metal is a serious candidate to explain these experimental observations. In addition, the new metallic phase in a dilute 2D electron gas can also be explained by the Bose metal phase if Cooper pairs form at the melting boundary of a 2D Wigner crystal as has been suggested previously [32].

Experimentally, measurements of the dc-conductivity are always made at finite frequency. As the frequency-dependent conductivity has a Lorentzian-type peak at $\omega = 0$ with a width of order $1/\tau$, the theoretical constraint on the experimental observation of the Bose metal is that $\omega \ll 1/\tau$. For a temperature of $0.1K$, the relaxation time, Eq. (7) is $10^{10}\exp(-m/T)\mathrm{sec}^{-1}$, where $m \ll T$. Typical experimental frequencies under which the dc-conductivity is measured are on the order of several Hz. Hence, the Bose metal can be observed provided that $T > 0.05m$. Typically, $T > 0.1m$. Consequently, there does not appear to be any experimental constraint regarding the frequency that prohibits the observation of the Bose metal phase.

What about dissipation? We have assumed in thus far that the only source of quasiparticle relaxation is the quartic term in the Ginzburg-Landau action. Meanwhile, other dissipative mechanisms are present in real experimental systems exhibiting the IST. For example, considering the dissipation-tuned [26,9,10] IST, Ohmic dissipation of the form $\eta|\omega_n|$ is always a relevant perturbation [18,29] arising generically from coupling to a lattice or an Ohmic resistor. For d-wave pairing, static disorder leads to Ohmic dissipation [30]. Provided that $\eta < m$, in the lowest approximation, the inverse relaxation time $1/\tau$ can be replaced by $1/\tau_{eff} = 1/\tau + \eta(k,T)$, where in the case of Ohmic dissipation $\eta(k,T) = \eta$, and our formula for the conductivity reads in the quantum disordered regime

$$\sigma(T) = 2\frac{(e^*)^2}{h}\frac{T}{\eta e^{m/T} + \pi T}. \qquad (17)$$

We see, that already for very small η the conductivity quickly starts to decrease exponentially with lowering temperature. This result is certainly intriguing and raises fundamental questions regarding the origin of the insulating state based on quantum fluctuations. We propose then that the more correct phase diagram for insulator-superconductor transitions is Fig. (1) in which it is clearly depicted that quantum fluctuations alone are insufficient to drive the insulating state. The finite temperature line starting at T_{KT} and terminating at g_c describes the standard Kosterlitz-Thouless transition. In the absence of dissipation, the T_{KT} line separates a superconductor and a Bose metal, not a superconductor and an insulator as had traditionally been thought. Systematic studies are needed in the presence of weak dissipation are needed to see if an insulator replaces the Bose metal phase at low temperatures.

ACKNOWLEDGMENTS

We thank the NSF grant No. DMR98-96134 for funding this work and A. Castro-Neto, S. Sachdev, E. Fradkin, A. Kapitulnik, and A. Yazdani for crucial discussions.

[1] S. Doniach, Phys. Rev. B **24**, 5063, (1981).
[2] M. P. A. Fisher, Phys. Rev. B **36**, 1917 (1987);
[3] X. G. Wen and A. Zee, Int. J. Mod. Phys. B **4**, 437 (1990).
[4] M.P.A. Fisher and G. Grinstein, Phys. Rev. Lett. **60**, 208, (1988).
[5] W. Zwerger, J. Low Temp. Phys. **72**, 291 (1988);ibid, Physica B **152**, 236 (1988).
[6] S. Chakravarty, S. Kivelson, G. T. Zimanyi, and B. I. Halperin, Phys. Rev. B **37**, 3283 (1988).
[7] A. Kampf and G. Schön, Phys. Rev. B **36**, 3651 (1987).
[8] V. Ambegoakar, U. Eckern, and G. Schön, Phys. Rev. Lett. **48**, 1745 (1982).
[9] K. Wagenblast, A. van Otterlo, G. Schön, and G. Zimanyi, Phys. Rev. Lett. **79**, 2730 (1997).
[10] K. Wagenblast, A. van Otterlo, G. Schön, and G. Zimanyi, Phys. Rev. Lett. **78**, 1779 (1997).
[11] A. van Otterlo, K.-H Wagenblast, R. Fazio, and G. Schön, Phys. Rev. B **48**, 3316 (1993).
[12] M. C. Cha, M. P. A. Fisher, S. M. Girvin, M. Wallin, and A. P. Young, Phys. Rev. B **44**, 6883 (1991).
[13] For a review see A. M. Goldman and N. Markovic, *Physics Today* **51** (11) 39 (1998).
[14] Y. Liu and A. M. Goldman, Mod. Phys. Lett. **8**, 277 (1994).

[15] A. Yazdani and A. Kapitulnik, Phys. Rev. Lett. **74**, 3037 (1995).
[16] N. Mason and A. Kapitulnik, Phys. Rev. Lett. **82**, 5341 (1999).
[17] K. Damle and S. Sachdev, Phys. Rev. B **56**, 8714 (1997).
[18] D. Dalidovich and P. Phillips, Phys. Rev. Lett. **84**, 737 (2000). Eqs. (5), (6), (8), (13), and (15) should be scaled by a factor of two.
[19] D. Das and S. Doniach, Phys. Rev. B **60** 1261 (1999).
[20] S. Sachdev, Phys. Rev. B **57**, 7157 (1998).
[21] A. Chubukov, S. Sachdev and J. Ye, Phys. Rev. B **49**, 11919 (1994).
[22] M. P. A. Fisher, and E. Fradkin, Nucl. Phys. B **251** [FS13] 457 (1985); E. Fradkin, Phys. Rev. B **33**, 3257 (1986); P. A. Lee, Phys. Rev. Lett. **71**, 1887 (1993).
[23] T. K. Kopec and J. V. Jose, Phys. Rev. Lett. **84**, 749 (2000).
[24] H. S. J. Van der Zant, et. al. Phys. Rev. B **54**, 10081 (1996).
[25] L.J. Geerligs, et. al. Phys. Rev. Lett. **63**, 326 (1989).
[26] A. J. Rimberg, et. al. Phys. Rev. Lett. **78**, 2632, (1997).
[27] T. K. Kopec and J. V. Jose, Phys. Rev. Lett. **84**, 749 (2000).
[28] P. W. Anderson, J. Phys. Chem. Solids **11**, 26 (1959).
[29] K. Voelker, cond-mat/9911473.
[30] I. F. Herbut, Phys. Rev. Lett. **85**, 1532 (2000).
[31] H. M. Jaeger, D. B. Haviland, B. G. Orr, and A. M. Goldman, Phys. Rev. B **40**, 182 (1989).
[32] P. Phillips, Y. Wan, I. Martin, S. Knysh, and D. Dalidovich, Nature **395**, 253 (1998).